Data changes between August 2017 evaluation run and January 2018 evaluation run.

New data received to ICBF database between 28th July 2017 and 24th November 2017.

| Trait | old | new | Extra | % Extra |
|-------------------------|------------|------------|---------|---------|
| calving difficulty | 13,117,186 | 13,417,747 | 300,561 | 2.2% |
| gestation | 4,217,626 | 4,262,712 | 45,086 | 1.1% |
| mortality | 18,063,938 | 18,372,695 | 308,757 | 1.7% |
| birth weight | 238,491 | 244,722 | 6,231 | 2.5% |
| weaning weight | 1,077,682 | 1,147,401 | 69,719 | 6.1% |
| yearling weight | 1,219,164 | 1,255,434 | 36,270 | 2.9% |
| Linear scores | 239,454 | 241,114 | 1,660 | 0.7% |
| weanling docility | 1,544,131 | 1,658,493 | 114,362 | 6.9% |
| weanling price | 1,069,162 | 1,117,318 | 48,156 | 4.3% |
| yearling price | 1,035,287 | 1,081,150 | 45,863 | 4.2% |
| Tully feed intake | 5,650 | 5,774 | 124 | 2.1% |
| carcass weight | 7,078,358 | 7,311,277 | 232,919 | 3.2% |
| carcass conformation | 7,078,358 | 7,311,277 | 232,919 | 3.2% |
| carcass fat | 7,078,358 | 7,311,277 | 232,919 | 3.2% |
| age 1st calving | 2,371,739 | 2,394,807 | 23,068 | 1.0% |
| calving interval | 4,627,019 | 4,647,900 | 20,881 | 0.4% |
| maternal wean wt | 926,039 | 984,932 | 58,893 | 6.0% |
| cow milk score | 2,361,776 | 2,572,720 | 210,944 | 8.2% |
| survival | 5,458,186 | 5,477,848 | 19,662 | 0.4% |
| cow docility | 2,322,632 | 2,350,435 | 27,803 | 1.2% |
| cow live weight | 972,942 | 1,011,949 | 39,007 | 3.9% |
| cull cow carcass wt | 1,740,189 | 1,795,311 | 55,122 | 3.1% |
| Genotypes | 1,141,581 | 1,163,750 | 22,169 | 1.9% |
| foreign ebvs calving | 25,022 | 28,285 | 3,263 | 11.5% |
| foreign ebvs weaning wt | 50,146 | 50,393 | 247 | 0.5% |
| foreign ebvs linear | 43,281 | 43,461 | 180 | 0.4% |
| foreign ebvs carcass | 29,929 | 30,127 | 198 | 0.7% |
| foreign ebvs milk | 50,955 | 51,203 | 248 | 0.5% |